

Gravitational Microlensing by the Globular Cluster Stars

by

Bohdan Paczyński

Princeton University Observatory, Princeton, NJ 08544-1001, USA

e-mail: bp@astro.princeton.edu

Received July 28, 1994

ABSTRACT

The microlensing of background stars by compact objects in globular clusters is analyzed. The main strength of the proposed search is the direct relationship between the lens mass and the time scale of the microlensing event. The main problem is the low overall rate of expected events which implies that a ground based search should last for about a decade to generate a non trivial number of events, like a few dozen. A space borne experiment could increase the rate by a large factor by increasing the number of stars which could be monitored thanks to the much better seeing. The clusters M22 with the galactic bulge background, and 47 Tuc with the SMC background, are the two examples of possible targets.

Key words: gravitational lensing - stars: clusters, low mass, brown dwarfs

The effect of gravitational microlensing of the LMC, SMC, and the galactic bulge stars by all intervening compact objects has been proposed as a way to detect low mass, dark objects, like brown dwarfs and even planets (Paczynski 1986, 1991, Griest et al. 1991). Recently, three groups reported the detection of the total of 17 microlensing event candidates: OGLE - 10 (Udalski et al. 1993, 1994a,b,c), MACHO - 5 (Alcock et al. 1993,

1994), and EROS - 2 (Auburg et al. 1993). The most thoroughly analyzed results were those of the OGLE (Udalski et al. 1994b, Paczyński et al. 1994), and these revealed a very unpleasant reality: the determination of the masses of lensing objects on the basis of the observed event time scales is very model dependent. In fact, there is so much uncertainty about the distribution of ordinary stars, not to mention brown dwarfs and more exotic objects, in our own galaxy that the mass determination is currently impossible. The aim of this paper is to present the conditions under which the relation between the time scale and the lens mass is very direct and reasonably unique.

Consider rich globular clusters, like 47 Tuc or M22, seen against the rich background of either SMC or the galactic bulge stars. We propose to consider the lensing events that are expected when any compact globular cluster object passes in front of a distant star, which acts as the source of light. A typical velocity dispersion in a globular cluster is $\sim 10 \text{ km s}^{-1}$, while a typical transverse velocity of a cluster as a whole is $\sim 100 \text{ km s}^{-1}$. Therefore, even if the lens cannot be directly detected its transverse velocity is well approximated by that of a cluster as a whole. The same is true for the distance to the lens. The lensed star is always directly visible, and in most cases it is likely to be at the well known distance, be it the SMC or the galactic bulge. As a reasonable first approximation one can take the distance to the source to be the same as that of the average system it belongs to. However, in principle its own distance is measurable, as it is directly observed.

The geometry as outlined above offers a unique opportunity to derive the lens mass when the event time scale is measured. The combined uncertainty due to the depth effects of the source system or the velocity dispersion in the cluster may generate $\sim 20\%$ uncertainty in the lens mass. The uncertainty with the currently reported lensing events is more like a factor of 3 or even 10, as in case of the LMC lensing events the lenses may be anywhere between the galactic disk and the LMC bar (Gould et al. 1994, Sahu 1994). Therefore, the lenses in globular clusters offer unprecedented accuracy of the mass determination.

There is a very good reason to determine the mass function in globular clusters, as there is observational indication that “some, (and perhaps all) clusters probably have

very steep IMF’s with the slope likely exceeding 2.5 (Salpeter value 1.3)” (Richer et al. 1991). The inference was based on the observed *luminosity* function and the theoretical mass-luminosity relation. The gravitational lensing offers the only direct determination of the mass function in the globular clusters which are seen against a rich background of reasonably bright stars.

Let us determine the number of events expected in a whole cluster and the relation between the event time scale and the lens mass following Paczyński (1991). For simplicity the mass distribution in a globular cluster is approximated here with a singular isothermal sphere truncated at 1/3 of its tidal radius as estimated with the King model (King 1966). A singular model is wrong within the core radius, but this is of no concern to us as no background stars can be detected within the core radius.

The column mass density of a singular isothermal sphere is given as

$$\Sigma \approx \frac{\sigma^2}{2Gr}, \quad (1)$$

where r is the projected distance from the cluster center. Let D_s and D_d be the distance from the observer to the source and the lens (deflector), respectively. The optical depth to microlensing can be calculated to be

$$\tau = \frac{4\pi G\Sigma}{c^2} D_d \left(1 - \frac{D_d}{D_s}\right) = \frac{\sigma^2}{c^2} \frac{2\pi}{\varphi} \left(1 - \frac{D_d}{D_s}\right), \quad (2)$$

where σ is the one dimensional (radial) velocity dispersion, and φ is the angular distance from the cluster center. If the sources are at very large distance, $D_d/D_s \ll 1$, then the correction term in the brackets is equal one, and the formula is even simpler:

$$\tau \approx 2.4 \times 10^{-5} \left(\frac{\sigma}{10 \text{ km s}^{-1}}\right)^2 \left(\frac{1'}{\varphi}\right). \quad D_d/D_s \ll 1. \quad (3)$$

This optical depth is much higher than typically expected towards the galactic bulge or the SMC, i.e. within a few arcminutes of the cluster center the lensing is likely to be dominated by the compact objects associated with the cluster itself. In any case the “background” due to the other galactic or LMC lenses can be readily determined observationally.

Following Paczyński (1991) we can also calculate the relation between the time scale of the microlensing event, $t_0 \equiv R_E/V$, where R_E is the Einstein ring radius, and V is the

transverse velocity in the observer-lens-source system. The time scale can be calculated as

$$t_0 [\text{days}] = 32.9 \left(\frac{M_d}{0.1 M_\odot} \right)^{1/2} \left(\frac{1 \text{ kpc}}{D_d} \right)^{1/2} \left(1 - \frac{D_d}{D_s} \right)^{1/2} \left(\frac{10 \text{ mas yr}^{-1}}{\dot{\varphi}} \right), \quad (4)$$

where M_d is the lens (deflector) mass, $\dot{\varphi}$ is the observed relative proper motion of the globular cluster with respect to the lensed star, and all other symbols have their usual meaning. As expected the time scale is shorter when the lens mass is smaller, the distance to the lens is larger, and its transverse angular velocity is larger.

Now we have to estimate the number of events that are expected in one year of continuous monitoring of the cluster background. The OGLE experience indicates (Udalski et al. 1992) that up to 400 stars can be measured per area $(1')^2$ in the sky while the seeing is $1''$ (FWHM). Naturally, the better the seeing the more stars can be measured in a dense stellar field. We adopt

$$n \approx 400 \left(\frac{1''}{FWHM} \right)^2 \quad \text{per } (1')^2, \quad (5)$$

where n is the number of measurable stars per $(1')^2$. The number of measurable background stars that are microlensed at any instant, i.e. which have their apparent brightness increased by at least a factor 1.34, can be calculated as

$$N = \int_{\varphi_1}^{\varphi_2} n \tau \, 2\pi \varphi \, d\varphi = 0.60 \left(\frac{1''}{FWHM} \right)^2 \left(\frac{\varphi_2 - \varphi_1}{10'} \right) \left(\frac{\sigma}{10 \text{ km s}^{-1}} \right)^2 \left(1 - \frac{D_d}{D_s} \right), \quad (6)$$

where φ_1 and φ_2 are the inner and the outer radius of the annulus around the cluster within which the search for microlensing is feasible. The rate of events per year can be calculated as

$$\begin{aligned} \Gamma [\text{yr}^{-1}] &= \frac{2N}{\pi} \frac{1 \text{ yr}}{t_0} = \\ &4.2 \times \left(\frac{\sigma}{10 \text{ km s}^{-1}} \right)^2 \left(\frac{0.1 M_\odot}{M_d} \right)^{1/2} \left(\frac{\varphi_2 - \varphi_1}{10'} \right) \left(\frac{\dot{\varphi}}{10 \text{ mas yr}^{-1}} \right) \times \\ &\quad \left(\frac{D_d}{1 \text{ kpc}} \right)^{1/2} \left(1 - \frac{D_d}{D_s} \right)^{1/2} \left(\frac{1''}{FWHM} \right)^2. \end{aligned} \quad (7)$$

The last two equations demonstrate the main problem with the idea presented in this paper: the expected rate of events is very small unless a substantial fraction of globular

cluster mass is in very low mass objects, or we can somehow reduce the FWHM of the seeing disk, for example with a space borne experiment. A traditional observing program from the ground should extend for about a decade to generate non-trivial results.

There is yet another problem. Our estimate of ~ 400 stars per square arcminute at the 1" seeing refers to the saturation limit: we do not expect to be able to measure more stars than that number. In case of M22 the background galactic bulge stars are bright enough that it might be relatively easy to reach the saturation limit with a 1 meter class telescope. The background of SMC stars in case of 47 Tuc is much fainter (cf. Fig. 4 of Hesser *et al.* 1987), and it may take a 4 meter class telescope to carry out the observing program.

Naturally, the range of angular distances from the cluster center, $\varphi_1 \leq \varphi \leq \varphi_2$, is limited by the crowding of the clusters stars at φ_1 , and the tidal radius of the clusters at φ_2 .

Acknowledgements. It is a great pleasure to acknowledge stimulating discussions with N. Reid and I. Thompson, and the hospitality of A. Omont and many other astronomers at the Institut d'Astrophysique in Paris where this paper has been written. This project was supported with the NSF grant AST92-16494.

REFERENCES

- Alcock, Ch. *et al.* 1993, *Nature*, **365**, 621.
- Alcock, Ch. *et al.* 1994, *Astrophys. J. Letters*, submitted.
- Aubourg, E. *et al.* 1993, *Nature*, **365**, 623.
- Gould, A., Miralda-Escudé, J. and Bahcall, J. N. 1994, *Astrophys. J.*, **423**, L105.
- Griest, K. *et al.* 1991, *Astrophys. J. Letters*, **372**, L79.
- Hesser, J. E., Harris, W. E., Vandenberg, D. A., Allwright, J. W. B., Shott, P., and Stetson, P. B. 1987, *Publ. Astron. Soc. Pacific*, **99**, 739.
- King, I. R. 1966, *Astron. J.*, **71**, 64.

- Paczynski, B. 1986, *Astrophys. J. Letters*, **304**, 1.
- Paczynski, B. 1991, *Astrophys. J. Letters*, **371**, L63.
- Paczynski, B., Stanek, K. Z., Udalski, A., Szymański, M., Kałużny, J., Kubiak, M., Mateo, M., & Krzemiński, W. 1994, *Astrophys. J. Letters*, submitted.
- Richer, H. B., Fahlman, G. G., Buonanno, R., Pecc, F. F., Searle, L., and Thompson, I. B. 1991, *Astrophys. J.*, **381**, 147.
- Sahu, K. 1994, *Nature*, ???, ...
- Udalski, A., Szymański, M., Kałużny, J., Kubiak, M., and Mateo, M. 1992, *Acta Astron.*, **42**, 253.
- Udalski, A., Szymański, M., Kałużny, J., Kubiak, M., Krzemiński, W., Mateo, M., Preston, G. W., and Paczynski, B. 1993, *Acta Astron.*, **43**, 289.
- Udalski, A., Szymański, M., Kałużny, J., Kubiak, M., Mateo, M., and Krzemiński, W. 1994a, *Astrophys. J. Letters*, **426**, L69.
- Udalski, A., Szymański, M., Stanek, K. Z., Kałużny, J., Kubiak, M., Mateo, M., Krzemiński, W., Paczynski, B., and Venkat, R. 1994b, *Acta Astron.*, **44**, 165.
- Udalski, A., Szymański, M., Mao, S., Di Stefano, R., Kałużny, J., Kubiak, M., Mateo, M., and Krzemiński, W. 1994c, *Astrophysical J. Letters*, submitted